

Perinatal Outcome In 1515 Cases Of Prolonged Second Stage

Getting the books **perinatal outcome in 1515 cases of prolonged second stage** now is not type of challenging means. You could not single-handedly going in the manner of books heap or library or borrowing from your associates to gain access to them. This is an agreed easy means to specifically acquire guide by on-line. This online publication perinatal outcome in 1515 cases of prolonged second stage can be one of the options to accompany you subsequently having other time.

It will not waste your time. resign yourself to me, the e-book will utterly announce you new issue to read. Just invest tiny era to right of entry this on-line pronouncement **perinatal outcome in 1515 cases of prolonged second stage** as skillfully as evaluation them wherever you are now.

Once you find something you're interested in, click on the book title and you'll be taken to that book's specific page. You can choose to read chapters within your browser (easiest) or print pages out for later.

Perinatal Outcome In 1515 Cases

Perinatal Outcome in 1515 Cases of Prolonged Second Stage of Labour in Nulliparous Women J Obstet Gynaecol Can . 2015 Jun;37(6):508-16. doi: 10.1016/s1701-2163(15)30227-9.

Perinatal Outcome in 1515 Cases of Prolonged Second Stage ...

Measures of perinatal outcome were determined in relation to the duration of the second stage of labour . Of the 1515 neonates, 77 (5.1%) had a five-minute Apgar score < 7, 36 (2.4%) were admitted to the NICU, and 10 (0.7%) had confirmed or suspected seizures.

Perinatal Outcome in 1515 Cases of Prolonged Second Stage ...

During the study period, 1515 women met the inclusion criteria. The majority of women (67%) delivered vaginally, after up to 10 hours in the second stage of labour The overall rate of Caesarean section was low (15.7%). The rate of permanent neurologic impairment was 2.6 per 1000 deliveries.

Perinatal Outcome in 1515 Cases of Prolonged Second Stage ...

Perinatal Outcome in 1515 Cases of Prolonged Second Stage of Labour in Nulliparous Women. Hunt JC(1), Menticoglou SM(1). Author information: (1)Department of Obstetrics, Gynecology and Reproductive Sciences, University of Manitoba, Winnipeg MB.

Perinatal Outcome in 1515 Cases of Prolonged Second Stage ...

Download Citation | Perinatal Outcome in 1515 Cases of Prolonged Second Stage of Labour in Nulliparous Women | To examine perinatal outcomes among nulliparous women with a second stage of labour ...

Perinatal Outcome in 1515 Cases of Prolonged Second Stage ...

1. J Perinat Med. 2014 Nov;42(6):769-75. doi: 10.1515/jpm-2014-0072. Adverse perinatal outcomes related to the delivery mode in women with monochorionic diamniotic twin pregnancies. Yamashita A, Ishii K, Taguchi T, Mabuchi A, Ota S, Sasahara J, Hayashi S, Mitsuda N.

Adverse perinatal outcomes related to the delivery mode in ...

The primary outcome was survival at 28 days after birth. RESULTS: Between 1 January 2005 and 31 December 2010, 1950 pregnant women carrying 2151 fetuses underwent a total of 2828 FEs. CHD was diagnosed in 152 fetuses (7.1%), and perinatal outcome was available in 120, among which 78 (65%) had consanguineous parents.

Perinatal outcome of congenital heart disease in a ...

The association between stillbirth in the first pregnancy and subsequent adverse perinatal outcomes. Am J Obstet Gynecol 2009;201:378.e1-6. 5. Heinonen S, Kirkinen P. Pregnancy outcome after previous stillbirth resulting from causes other than maternal conditions and fetal abnormalities. Birth 2000;27:33-7. 6.

Stillbirth and subsequent pregnancy outcome - a cohort ...

Van Vliet et al. recently published, that in women with threatened preterm birth, 48 h of tocolysis with nifedipine or atosiban resulted in similar perinatal outcomes . However, more cases of intrauterine fetal demise (IUID) were registered in the nifidipine group (5% vs. 2% in the atosiban group; RR 2.20, 95% CI 0.91-5.33, n.s.).

Mid-trimester preterm premature rupture of membranes ...

Angela Yulia, Oana Chelemen, Ruchi Bhutani, Ciaran Hutchinson, Samantha Levine, Dia Kamel, Ruwan Wimalasundera, Large placental chorioangioma: a potential effective in-utero treatment modality for radio frequency ablation, Case Reports in Perinatal Medicine, 10.1515/crpm-2018-0040, 8, 2, (2019).

Color Doppler imaging of placental masses: differential ...

The main objective of this retrospective analysis in a large tertiary center was the clinical outcome of prenatally diagnosed cardiac rhabdomyomas as well as the identification of factors influencing fetal prognosis. Methods. A total of 45 cases of fetuses with prenatally suspected rhabdomyoma and their clinical outcome were analyzed ...

Clinical outcome of prenatally suspected cardiac ...

In selected cases, amniodrainage allows continuation of the pregnancy with improving perinatal outcome. Fetuses who develop hydrops are at the highest r ... Placental chorioangioma is associated with an increased risk of pregnancy complications, the most common being polyhydramnios and preterm delivery.

Perinatal Outcome After Prenatal Diagnosis of Placental ...

Perinatal outcomes with isolated oligohydramnios at term pregnancy. Naveiro-Fuentes M, Puertas Prieto A, Ruíz RS, Carrillo Badillo MP, Ventoso FM, Gallo Vallejo JL. AIMS: To compare the outcomes of term gestations with oligohydramnios in the absence of other underlying disorders and term gestations with normal amniotic fluid.

Perinatal outcomes with isolated oligohydramnios at term ...

Abstract. Objective: Gitelman syndrome (GS) is a rare renal disease, originating from a defect in the Na-Cl co-transporter in the distal tubule, which causes hypokalemia, hypomagnesemia, hypocalciuria, metabolic alkalosis and low-normal blood pressure. Mild hypokalemia of pregnancy is physiological. Fatigue, nausea, vomiting, polyuria and low blood pressures are also common complaints ...

Gitelman syndrome during pregnancy - from diagnosis to ...

Background and aim. To compare perinatal outcome in groups of planned vaginal breech delivery, elective cesarean section with the fetus in breech presentation, and planned vaginal delivery with the fetus in cephalic presentation in a university hospital with a tradition of managing breech deliveries by the vaginal route. Methods. A cohort study from a 7-year period 1995-2002, including 590 ...

Good perinatal outcome in selective vaginal breech ...

Objective: To compare the results of maternal and perinatal pregnancies of obese women after bariatric surgery. Methods: A retrospective cross-sectional study was carried out on 63 women who had undergone bariatric surgery and 73 obese women (control). Demographic data, the characteristics of the bariatric surgery, and the maternal and perinatal results were evaluated.

Maternal and Perinatal Outcomes After Bariatric Surgery: A ...

Objective: To evaluate the effect of coronavirus disease 2019 (COVID-19) on maternal, perinatal and neonatal outcome by performing a systematic review of available published literature on pregnancies affected by COVID-19. Methods: We performed a systematic review to evaluate the effect of COVID-19 on pregnancy, perinatal and neonatal outcome.

Effect of coronavirus disease 2019 (COVID-19) on maternal ...

Christos Parthenis, Vasiliki Michalitsi, Chrysanthi Chlapoutaki, Athena P. Souka, Prenatal diagnosis of fetal hemivertebra in the first trimester, Case Reports in Perinatal Medicine, 10.1515/crpm-2018-0050, 0, 0, (2019).

Fetal hemivertebra: associations and perinatal outcome ...

Judith Altmann, Verena Kiver, Wolfgang Henrich, Alexander Weichert, Clinical outcome of prenatally suspected cardiac rhabdomyomas of the fetus, Journal of Perinatal Medicine, 10.1515/jpm-2019-0246, 0, 0, (2019).

Copyright code: d41d8cd98f00b204e9800998ecf8427e.